

High Fluoride Concentration in Ground Water in Parts of Chhattisgarh – A Review

Meena Chakraborty¹, Dr. Madhurima Pandey*², Dr. Piyush Kant Pandey³

¹Govt. College Bori, Durg, Chhattisgarh

²Bhilai Institute of Technology Durg, Chhattisgarh

³Bhilai Institute of Technology Raipur, Chhattisgarh

¹E mail: chakrabortymeena@gmail.com

*²E mail: drmadhurima_pandey@yahoo.com

³E mail: drpiyush_pandey@yahoo.com

Abstract

Groundwater quality problems have emerged in many geographical areas due to natural environmental processes and human intervention in the geosystems. Among the water quality parameters, fluoride ion exhibits unique properties as its concentration in optimum dose in drinking water is advantageous to health and if the concentration exceeds the limit, this affects the health. High fluoride concentration in the ground water and surface water in many parts of the world is a cause of great concern. India has acute public-health problems induced by utilization of groundwater as a source of drinking water. The main source of fluoride in ground water is fluoride-bearing rocks This paper will focus on fluoride concentration and its relationship to water-quality parameters and its impacts on humans through groundwater resources in different parts of Chhattisgarh.

Keywords: Groundwater, Geosystem, Contamination, fluoride-bearing rocks

1.Introduction

Fluoride in small concentration (upto 0.6 mg/L) in drining water is essential for good dental health but higher concentration of fluoride (more than 1.5 mg/L) causes dental and skeletal fluorosis [1, 2]. High fluoride concentration in the ground water in many parts of the world is a cause of great concern. Ground water is considered to be in purest form, but as it comes in contact with rocks due to dissolution of minerals it becomes polluted. The main source of fluoride in ground water is fluoride-bearing rocks such as fluorspar, fluorite, cryolite, fluorapatite and hydroxylapatite [3,4]. Also the content in ground water is a function of many factors such as availability and solubility of fluoride minerals, velocity of flowing water, pH, temperature, concentrations of calcium and bicarbonate ions in water [1,5,6].

It is estimated that around 200 million people among from 25 nations the world over are under the dreadful fate of fluorosis. The higher concentration of fluoride in ground water associated with igneous & metaphoric rocks such as granites and gneisses have been reported from China, India, Pakistan [7,8,9]. In Indian the disease is endemic in about 275 districts of 20 states [10]. The most seriously affected

areas are Andhra Pradesh, Punjab, Haryana, Rajasthan, Gujrat, Tamilnadu & Utter Pradesh [11,12,13,14].

Chhattisgarh is a newly formed state of India (1st November 2000) after partinoing Madhaya Pradesh. A resource-rich state, it is a source of electricity and steel for the country, accounting for 15% of the total steel produced [15]. Chhattisgarh is one of the fastest-developing states in India [16]. Fluoride is a new threat determined for the population of Chhattisgarh. In Chhattisgarh endemic pockets of fluoride were found in Durg , Bastar, Kanker, Surguja, Surajpur, Balrampur, Balod and Korba districts [10]. This paper will focus on high fluoride concentration in some parts of Chhattisgarh.

2.Source of Fluoride contamination

2.1 Geogenic source

Fluoride in ground water is mainly a geogenic contamination, when ground water comes in contact with rocks containing fluoride bearing minerals, then dissolution of these minerals increases fluoride concentration in ground water. The ultimate concentration of fluoride in groundwater largely depends on reaction times with aquifer minerals. High fluoride concentrations can be built up in groundwaters which have long residence times in the aquifers. Such groundwaters are usually associated with deep aquifer systems and a slow groundwater movement [5]. The common fluorine bearing mineral and their formula is shown in Table 1.

Table 1: Chemical composition of fluoride bearing minerals

Name of mineral	Chemical Formula
Topaz	$Al_2(F,OH)_2SiO_4$
Fluoroapatite	$Ca_{10}(PO_4)_6F_2$
Fluorite	CaF_{2s}
Villiaumite	NaF
Sellaite	MgF_2
Cryolite	Na_3AlF_6
Wagnerite	Mg_2PO_4F
Mica	$AB_{2-3}(X,Si)_4O_{10}(O,F,OH)_2$
Vesuvianite	Ca,Mg Silicates with F
Tourmaline	Borosilicate with F

2.2 Chemical compositon of ground water

High-fluoride groundwaters are mainly associated with a sodium-bicarbonate water type and relatively low calcium and magnesium concentrations. Such water types usually have high pH values. Information on chemical composition of groundwater can be used as an (proxy) indicator of potential fluoride problems [1].

If calcium is present in higher concentration it is most effective in reducing the fluoride concentration. A strong negative Correlation has been observed between Ca and F in the ground waters that contain Ca in excess of that required for the solubility of fluoride minerals.

The positive correlation between F⁻ and Na⁺ & silica indicates the source of F⁻ in ground water to be from weathering of silicate minerals. The positive correlation between F and Na⁺ indicates dissolution of feldspar from feldspathic sandstones.

Since the increase in Na⁺ concentration increases the F⁻ bearing minerals, the geochemical process leading to increase in conc. of Na⁺ & decrease in Ca²⁺ plays an important role in F⁻ enrichment in ground water.

2.3 Anthrapogenig activities

Many industries such as aluminum industry, steel production plants, superphosphate plants, ceramic factories, coal-burning power plants, brickworks, glassworks, and oil refineries discharge effluents containing fluoride in them to water bodies.

3. Impact of fluoride on health

3.1 Dental fluorosis

Dental fluorosis is minerlization of tooth enamel caused by ingestion of high concentration of fluoride during enamel formation [17,18]. The severity of the condition is dependent on the dose, duration, and age of the individual during the exposure. There are three stages of dental fluorosis. In the "very mild" stage small, opaque, white areas scattered irregularly over the tooth, then in the "mild" form mottled patches developes on half of the surface area of the teeth. But in moderate form, entire teeth get mottled and teeth may be ground down and brown stains occur on the teeth. Severe fluorosis is characterized by brown discoloration and discrete or confluent pitting; brown stains are widespread and teeth often present a corroded-looking appearance [17]. People with fluorosis are relatively resistant to dental caries [19]. Dental fluorosis occurs due to conversion of hydroxyapatite into fluorohydroxyapatite.

3.2 Skeletal fluorosis

Skeletal fluorosis is a bone disease. It happens due to excessive accumulation of fluoride in the bones. Presence of high concentration of fluoride in the body causes harding of bones and they become less elastic Therefore such bones are more keen to fracture. Other symptoms include thickening of the bones and accumulation of fluoride in bone tissues, which makes the joint mobility impossible. Ligaments and cartilage can become ossified [20]. In advanced stages, skeletal fluorosis causes painful damage to bones and joints. Severity of the desease can lead to neurological defects or compression of spinal cord.

3.3 Other diseases

High fluoride concentration in body causes gastrointestinal problems, urinary tract malfunctioning, nausea, abdominal pain, tingling sensation in fingers and toes, reduced immunity, repeated abortions or still births, male sterility, etc [21]and also affect the functioning of kidneys [22].

Several cases of fluorosis in human beings and animals are also reported in different parts of Chhattisgarh (Table 2).

Table 2: Fluorosis reported in parts of Chhattisgarh

Study area	Fluorosis reported	Reference
Raigarh	Dental & skeletal fluorosis in humans	[23]
Rajnandgaon	Skin, dental and skeleton fluorosis in humans and domestic animals	[24]
Balod	Skeletal fluorosis in humans	[25]
Surajpur	fluorosis in humans	[26]
Kanker	Dental & skeletal fluorosis in humans	[10]
Durg	Dental & skeletal fluorosis in livestock population	[27]

4. Fluoride in parts of Chhattisgarh

In this paper fluoride contaminated areas of six district of chhattisgarh were considered i.e. Raigarh, Rajnandgaon, Kanker, Surajpur, balod and Durg. Description of these areas are given in breif in Table 3.

Table 3: Description of parts of Chhattisgarh with high fluoride concentration

Study area	Location Description	Geology	Reference
Tamnara block, Raigarh	lat. 22°05'N and 22°15'N, and long. 83°20'E and 83°30'E, covering about 240 sq. km area.	The area is covered by the rocks of Barakar, Barren measure, Raniganj and Kamthi formations of Gondwana Supergroup, consisting of a thick sequence of sandstones, shales, carbonaceous shales, clays and coal seams.	[23]
Kourikasa in Ambargarh Chouki block, Rajnandgaon	20°43'9''N and 80°44'7''E	Rocky basement covered by alluvium, colluvium, and soil. Tube wells tap groundwater in the fracture system in bedrock.	[24]
Dongargarh, RJN	21° 06' N 81° 02' E	-	[4]
Balod	area of the district is ≈4000 km ² with population of 1.3 million	-	[25]
Surajpur	area of 16034.4 Sq.kms with 54 percent of tribal population	Archean, Gondwanas, Lametas and Deccan trap group of rocks overlain by sub Recent to Recent alluvial sediments.	[26]
Domarpani village is located in a Narharpur block Kanker	Domarpani village is located in a Narharpur block of Kanker district with population 1593	-	[10]
Durg	Bemetara, Saja, Navagarh, Berla, Patan, Dhamdha		[27]

5. Methods of analysis of fluoride in water samples

Fluoride in water samples is generally analysed by colorimetric method or by potentiometric ion selective method (ISE). ISE method gives accurate result but equipment is costly whereas colorimetric method is less time consuming and cost effective [28]. Fluoride Ion Selective Electrode consists of a single crystal of lanthanum fluoride as the membrane. Only fluoride ions are mobile to this membrane. When it comes in contact with fluoride solution, a potential develops across the membrane which is directly proportional to the fluoride ions in the solution. In SPADNS method fluoride reacts with certain zirconium dyes to form a colourless complex and another dye. The dye becomes progressively lighter as fluoride concentration increases. Absorbance is measured at 570 nm and fluoride values for the samples were determined by plotting calibration curves. Whereas in case of alazarine complex method fluoride decolorizes the zirconium alazarine complex and decolorization is proportional to fluoride concentration in water. Description of these methods are given in Table 4.

Table 4 : Analytical methods for determination of fluoride in water samples

Analytical method	Preparation	Detection limit	Reference
Colorimetry	Dilute sample; add barium chloride; complex with zirconiumxylenol orange for color development	2,000 µg/L	[29]
Colorimetry	Sample added to sulfuric acid and distilled to remove interferences; distilled sample treated with SPADNS reagent; color loss resulting from reaction of reagent with fluoride is determined at 570 nm and concentration read off standard curve	0.10 mg/L	[30]
Colorimetry	Bellack distillationa , after which fluoride ion reacts with the red cerous chelate of alizarin complexone in an autoanalyzer	0.050 mg/L	[31]
Ion selective electrode	Mix sample and standard 1:1 with TISAB (for soluble fluorides)	0.500 mg/L	[32]
Ion selective electrode	No sample treatment required	0.100 mg/L	[32]

Analysis method adopted for determination of fluoride in water samples in the areas discussed in this paper are shown in Table 4.

Table 4: Analysis method adopted for fluoride in water sample in selected areas

Study area	No. of samples	Method used for F ⁻ determination	Reference
Tamnar block, Raigarh	164	ion chromatograph (Metrohm, 861, Advanced Compact IC)	[23]
Kourikasa in Ambagarh Chouki block, Rajnandgaon	20	Orion Ion Analyzer-720 equipped with the ion-selective electrode using the buffer (TISAB-III)	[24]
Dongargarh, RJN	15	SPADNS	[4]
Balod	50	Metrohm ion meter-781 using the CDTA buffer	[25]
Surajpur	30	SPADNS	[26]
Domarpani village is located in a Narharpur block Kanker	41	Analysis was done by Public Health Engineering laboratory at district headquarter (Kanker)	[10]
Durg	168	Digital Ion Analyser equipped with a fluoride-selective electrode (Orion Research, Model 701A, Massachusetts) with buffer (TISAB) II	[27]

6. Fluoride concentration water iledcted areas of Chhattiisgarh

6.1 Fluoride in Raigarh district

In Raigarh district the fluoridated area is a part of the Pahaj River watershed in Tamnar Block [23], where fluoride in water was first reported in 2004 by the Public Health Engineering Department of the State Government. Beg et al. reported that F⁻ concentration in groundwater consistently exceeded the desirable and maximum permissible limits in three villages, viz. Muragaon, Pata and Saraitola. high F⁻ concentration in groundwater mainly occurred in wells tapping the aquifers in Barakar Formation, which has a litho-assemblage of feldspathic sandstone/shale/coal. The positive correlation of F⁻ with Na⁺ and SiO₂, the groundwater type, increase in Na⁺ : Ca²⁺ values with increase in F⁻ concentration, presence of Li⁺ in the high-F⁻ zone, absence of PO₃⁴⁻, and the results of mineralogical and petrographic analyses indicate that F⁻ in groundwater is geogenic.

6.2 Fluoride in Rajnandgaon

In rajnandgaon district the ground water were collected from fifteen bore wells from different parts of the three blocks such as Dongargarh, Dongargoan and Rajnanadgaon in the month of January 2015 [4] out of which two sites of Dongargarh has fluoride more than permissible limit in water. In their paper of Noel & Rajwade, 2015 [4] physicochemical parameters pH, Electrical conductivity (EC) and total dissolved solid (TDS) and total hardness (TH) was determined but they have not correlated these parameters with presence of fluoride in water samples.

In another stusy made by Patel et al., 2015 [24] water samples were collected from Kourikasa area of Ambagarh Chouki, Rajnandgaon (RJN) district in 2010-11. In this area average concentration of fluoride in water was 14.1 ± 3.1 which is several times more than the permissible limit. By statistical analysis they

found that fluoride contamination of the groundwater seems to be related to the tube wells installed at shallow depth and, thus, to enrichment of shallow ground water by evaporation.

6.3 Fluoride in Surajpur

Upadhyay & Komal Kumari, 2013 [26] collected six water samples from each of five villages namely Baraul, Ramtirath, Banapatti, Barwahi and Fatehpur from Surajpur district for fluoride analysis. Out of these six samples three were surface and three were ground water samples.

In the study made by Upadhyay & Komal Kumari, 2013 [26], all the samples collected from five villages of Prem Nagar block of Surajpur showed fluoride concentration more than prescribed permissible limit. Water quality parameters such pH, temperature, conductivity, dissolved oxygen T.D.S. were analysed but they didn't report any correlation of these parameters with fluoride. Therefore source of fluoride is not detectable from this study.

6.4 Fluoride in Balod

Yadav et al., 2016 [25] reported fluoride contamination in ground water in Balod district. Samples were collected from Balod, Gurur, Gunderdehi, Dondi Lohara and Dondi block of Balod district. The physical parameters *i.e.* temperature (T), pH, dissolved oxygen (DO), reduction potential (RP) and electrical conductivity (EC) with Fe in samples were measured. Iron, fluoride, chloride and calcium were found more than the permissible limit in samples which made them not suitable for drinking purpose. This paper also reported skeletal fluorosis in Gunderdehi and Balod blocks. The source of fluoride in water in this area was found to be geogenic. Presences of fluoride bearing rocks are mainly responsible for the contamination. Mining and agricultural activities also contributes in enhancement of fluoride contamination in water sources of area.

6.5 Fluoride in Kanker

A survey report of fluorosis mapping was prepared by Dr. Gitte [10] of Domarpani village in Narharpur block of Kanker district. The objective of the survey was to determine the concentration of fluoride in drinking water and to find out the deformity caused by fluorosis in population of the area. Survey reported that dental fluorosis was common in teen age group whereas skeletal fluorosis was found in male and female above 45 years of age. The mean fluoride concentration in the ground water in the surveyed area was 2.19 and 98% of surveyed population is suffering from fluorosis there.

6.6 Fluoride in Durg

Giri et al., 2013 [27] made a cross sectional survey in 12 blocks of Durg district to determine the fluoride concentration in water. For the purpose, 168 water samples (7 groundwater samples and 7 surface water samples from each block) were collected. This paper considered that high concentration of fluoride in water is geogenic. The weathering and leaching process helps to increase fluoride contamination. Along with this industrial development of the area also contributes in fluoride contamination specially phosphate fertilizer factory and the brick kilns.

Fluoride concentration in areas discussed above and cause of fluoride in water samples there is summarized in Table 5.

Table 5: Fluoride concentration in parts of Chhattisgarh

Location	Min.	Max	Correlaton		Inference	Referenc e
			Positive	Negative		
Tamnar block, Raigarh	0.09	8.88	Na ⁺ and SiO ₂	ca ²⁺ , Mg ²⁺ , HCO ₃ ⁻ , alkalinity and total hardness (TH)	The F ⁻ rich groundwater in the study area is associated with Na-Ca-Mg-HCO ₃ and Na-Mg-Ca-HCO ₃ types of water	[23]
Ambagarh Chouki, RJN	3.7	27			Enrichment of shallow ground water by evaporation	[24]
Dongargarh, RJN	0.25	1.7	-	-	-	[4]
Premnagar Block, Surajpur	0.7	1.2	-	-	Geogenic contamination	[26]
Domarpani village in Narharpur block anker	0.5	2.86	-	-	-	[10]
Gunderdehi block, Balod	1.5	14	NO ₃ ⁻ , Na ⁺ , K ⁺ , Mg ²⁺ , Ca ²⁺ , NH ₄ ⁺	Cl ⁻ , SO ₄ ²⁻ , Fe	Positive correlation with the NH ₄ ⁺ , Na ⁺ , K ⁺ and Mg ²⁺ ions, indicating existence of F ⁻ in the geo media as Barberiite, bararite, ferruccite, sellatite, cryolite, hieratite, etc. Other ions i.e. Cl ⁻ , SO ₄ ²⁻ , Na ⁺ , K ⁺ , Mg ²⁺ and Ca ²⁺ were partially correlated, showing origin from multiple sources i.e. geogenic, mining, agriculture, etc.	[25]
Durg	0.2	13.2	-	-	geogenic contamination, along with contamination from phosphate fertilizer factory and the brick kilns	[27]

7. Necessary steps to be taken to prevent fluorosis

7.1 Use of alternative sources of water

Water is the main source of fluoride intake in body. Fluoride is mainly geogenic contamination and ground water is the main source of drinking in Chhattisgarh. Therefore use of alternative of ground water such as surface water or rain water should be encouraged in villages.

Moreover fluoride is not evenly distributed in ground water. Its distribution depends on geological structure and depth of bore wells. Hence digging of new bore well in area of low fluoride concentration can provide safe drinking water tp population.

7.2 Development of methods of removal of fluoride from water

Various methods are available for defluoridation of water such as chemical precipitation, adsorption, ionic separation etc. Each and every method has its own advantages and disadvantages. As per the condition and requirement of a fluoride contaminated site proper method can be adopted for defluoridation.

7.3 Nutrition:

Proper measures can be taken to provide nutritional supplements to people suffering from fluorosis. As calcium has negative correlation with fluoride, intake of calcium supplements along with vitamin C, iron, antioxidants such as milk, curd, green leafy vegetables, fruits is helpful for fluorosis patients.

7.4 Health Education and public awareness:

Awareness campaign should be run by government and health workers to provide information on fluorosis, importance of drinking safe water and about healthy diet are important for prevention and control of fluorosis. High fluoride containing products such as Supari, tobacco, black rock salt, red rock, toothpaste, mouth wash etc. should be avoided.

8. Conclusion

Fluoride in ground water is primarily derived from decomposition, dissociation and dissolution of F^- bearing minerals, and occasionally from anthropogenic activities such as use of phosphatic fertilizers which have F^- as an impurity. Ascertaining the source of F^- in groundwater is essential for taking up mitigation measure. All the reported areas discussed in this paper have high fluoride concentration in water. Some papers not only identified the fluoride contaminated areas but also correlate it with geological structure and chemical composition of water. Rock structure of area and correlation matrix of fluoride with other physicochemical parameters helps to identify fluoride contamination in water sources, which in turns help to adopt proper preventive measure in contaminated area. Therefore identification of fluoride in water, cause of fluoride contamination and development of removal method is essential to fight back the problem of fluorosis.

References

- [1] B. K. Handa, "Geochemistry and Genesis of Fluoride Containing Groundwater in India", *Ground Water*, vol. 13, (1975), pp. 275-281.
- [2] L. Ripa and C. Clark, "Water Fluoridation. In: Primary Preventive Dentistry, Editorial El Manual Moderno, Mexico, (2001), pp. 127-160.
- [3] B.S., Sharma, Jyoti Agrawal and A. K. Gupta, "Emerging Challenge: Fluoride Contamination in Groundwater in Agra District, Uttar Pradesh", *Asian Journal of Experimental Biological Sciences.*, vol. 2, (2011), pp. 131-134.
- [4] Megha Nooel & Ram Prakash Rajwade, "Study of Fluoride Concentration in Groundwater of Rajnandgaon District, Chhattisgarh, India", *International Journal of Research in Engineering Science and Technologies*, vol. 1, no. 4, (2015), pp. 50-52.
- [5] P. F. Hudak and S. Sanmanee, "Spatial Patterns of Nitrate, Chloride, Sulfate and Fluoride Concentrations in the Woodbine Aquifer of North-Central Texas", *Environmental Monitoring and Assessment*, vol. 82, (2003), pp. 311-320.

- [6] R. Brunt, L. Vasak and J. Griffioen, "Fluoride in Groundwater: Probability of Occurrence of Excessive Concentration on Global Scale. Report SP 2004-2", International Groundwater Resources Assessment Centre (IGRAC).
- [7] Q. Xiang, Y. Liang, L. Chen, C. Wang, B. Chen, X. Chen and M. Zhou, "Effect of Fluoride in Drinking Water on Children's Intelligence", *Fluoride*, vol. 36, (2003), pp. 84-94.
- [8] N. Srinivasa Rao, "The Occurrence and Behaviour of Fluoride in the Groundwater of the Lower Vamsadhara River basin, India". *Hydrological Sciences*, vol. 42, no. 6, (1997), pp. 877-892.
- [9] Farooqi. A., Masuda. H., Kusakabe. M., Naseem. M. and Firdous. N., "Distribution of Highly Arsenic and Fluoride Contaminated Groundwater from east Punjab, Pakistan, and the Controlling Role of Anthropogenic Pollutants in the Natural Hydrological Cycle", *Geochemical Journal*, vol. 41, (2007), pp. 213-234.
- [10] S. Gitte, Fluorosis baseline survey of village Domarpani, District Kanker (Tribal) Chhattisgarh state, survey report (2013).
- [11] S.D. Dhimana and Ashok K. Keshari, Hydrogeochemical evaluation of high-fluoride groundwaters: a case study from Mehsana District, Gujarat, India, *Hydrological Sciences Journal*, vol. 51, no. 6, (2006), pp. 1149-1162.
- [12] A.K. Yadav and Parveen Khan, Fluoride and Fluorosis Status in Groundwater of Todaraisingh Area of District Tonk (Rajasthan, India): A Case Study, *International Journal of Chemical, Environmental and Pharmaceutical Research*, vol. 1, no. 1, (2010), pp. 6-11.
- [13] Nagaraju Arveti · M. R. S. Sarma · J. A. Aitkenhead-Peterson · K. Sunil, Fluoride incidence in groundwater: a case study from Talupula, Andhra Pradesh, India, *Environ Monit Assess*, DOI 10.1007/s10661-010-1345-3.
- [14] V. Veeraputhiran and G. Alagumuthu., "A report on Fluoride distribution in drinking Water", *International Journal of Environmental Sciences*, vol. 1, no. 4, (2010), pp. 49-54.
- [15] "Chhattisgarh State – Power Hub". Archived from the original on 20 November 2010. Retrieved 22 July 2011.
- [16] "Chhattisgarh -Steel". Archived from the original on 7 July 2011. Retrieved 22 July 2011.
- [17] J.A. Dean, McDonald and Avery's Dentistry for the Child and Adolescent (10th ed.). Elsevier Health Sciences. ((2015)), pp. 132. ISBN 978-0-323-28746-3.
- [18] M.C. Wong, A. M. Glenny, B.W. Tsang, E.C. Lo, H.V. Worthington, V.C. Marinho, "Topical fluoride as a cause of dental fluorosis in children". *The Cochrane Database of Systematic Reviews* (1): (2010), CD007693. doi:10.1002/14651858.CD007693.pub2. PMID 20091645.
- [19] B.W. Neville, A.C. Chi, D.D. Damm and C.M. Allen, *Oral and Maxillofacial Pathology* (4th ed.). Elsevier Health Sciences, (2015), pp. 52–54. ISBN 978-1-4557-7052-6.
- [20] L.V. Kalia, L. Lee, S.K. Kalia, F. Pirouzmand, M.J. Rapoport, R.I. Aviv, D. Mozeg and S.P. Symons, "Thoracic myelopathy from coincident fluorosis and epidural lipomatosis", *Canadian Journal of Neurological Sciences*, vol. 37, no. 2, (2010), pp. 276–278.
- [21] Meenakshi. and R. C. Maheshwari, "Fluoride in Drinking Water and its Removal", *Journal of Hazardous Materials*, vol. 137, (2006), pp. 456–463.
- [22] K. Brindha and L. Elango, "Fluoride in Groundwater: Causes, Implications and Mitigation Measures", *Fluoride Properties, Applications and Environmental Management*, (2011), pp. 111-136. ISBN:978-1-61209-393-2.
- [23] M. K. Beg, S. K. Srivastav, E. J. M. Carranza and J. B. de Smeth, "High fluoride incidence in groundwater and its potential health effects in parts of Raigarh District, Chhattisgarh, India", *Current Science*, vol. 100, (2011), pp.750 – 754.
- [24] Khageshwar Singh Patel, Bharat Lal Sahu, Nohar Singh Dahariya, Amarpreet Bhatia, Raj Kishore Patel, Laurent Matini, Ondra Sracek and Prosun Bhattacharya, "Groundwater arsenic and fluoride in Rajnandgaon District, Chhattisgarh, northeastern India", *Appl Water Sci*, (2015), DOI 10.1007/s13201-015-0355-2
- [25] Ankit Yadav, Yaman Kumar Sahu, Keshaw Prakash Rajhans, Pravin Kumar Sahu, Suryakant Chakradhari, Bharat Lal Sahu, Shobhana Ramteke, Khageshwar Singh Patel, "Fluoride Contamination of Groundwater and Skeleton Fluorosis in Central India", *Journal of Environmental Protection*, vol. 7, (2016), pp. 784-792.

- [26] Manish Upadhyay and Komal Kumari, “Determination of Fluoride Around In Surajpur District Chhattisgarh, India”, J. Atoms and Molecules, vol. 3, no. 1, (2013), pp./ 437–447
- [27] D. K. Giri, R. C. Ghosh, S. Dey, M. Mondal, Deepak Kumar Kashyap, Govina Dewanagan, “Incidence of hydrofluorosis and its adverse effects on animal health in Durg district, Chhattisgarh”, Current Science, vol. 105, no. 11, (2013), pp. 1477-1479.
- [28] Burton L. Edelstein, DDS, MPH David Cottrel, DDS David O’Sullivan, BS Norman Tinanoff, DDS, “Comparison of colorimeter and electrode analysis of water fluoride”, MS Pediatric Dentistry: January/February, vol. 14, no. 1, (1992), pp. 47-49.
- [29] Anthony G. Macejunas, Spectrophotometric Determination of Fluoride Using Zirconium-Xylenol Orange, American Water Works Association, vol 61, no. 6, (1969), pp.311-313.
- [30] Method 8029 FLUORIDE (0 to 2.00 mg/L F) for water, wastewater and seawater
- [31] N. Manivasakam , Physico-chemical Examination of Water, Sewage and Industrial Effluents. 3rd Edn. (1996), Pragati Prakashan, Coimbatore.
- [32] Method 9214 Potentiometric Determination of Fluoride In Aqueous Samples With Ion-Selective Electrode