

# INFLUENCE OF PLANT GROWTH REGULATORS ON GROWTH CHARACTERS OF DOLICHOS BEAN (*Lablab purpureus* (L) Sweet)

P. JAISANKAR\*, and K.MANIVANNAN,

*Department of Horticulture, Faculty of Agriculture, Annamalai University,  
Annamalai nagar, Tamilnadu.*

## ABSTRACT

A field experiment was conducted in 2017 at farmer's field in varagoorpatai village, Cuddalore, Tamilnadu to study the influence of plant growth regulators on growth characters of Dolichos bean (*Lablab purpureus* (L) Sweet). The experiment was laid out in Randomized Block Design with three replications and eleven treatments. The treatment combination consisted of an inorganic fertilizer (DAP) and Plant bio regulators (Panchagavya, Vermiwash, Effective Micro organism and Seaweed extract). The treatment details viz., T<sub>1</sub> - RDF (control), T<sub>2</sub> - INM (Without foliar application), T<sub>3</sub> - INM + DAP @ 2%, T<sub>4</sub> - INM + Panchagavya @ 3%, T<sub>5</sub> - INM + Panchagavya @ 4%, T<sub>6</sub> - INM + Vermiwash @ 1:5 dilution, T<sub>7</sub> - INM + Vermiwash @ 1:7.5 dilution, T<sub>8</sub> - INM + Effective Micro organism @ 1:1000 dilution, T<sub>9</sub> - INM + Effective Micro organism @ 2.5:1000 dilution, T<sub>10</sub> - INM + Seaweed extract @ 2.5% foliar spray, T<sub>11</sub> - INM + Seaweed extract @ 5%. The foliar sprays were applied at 30, 60 and 75 DAS. Studies revealed that the growth characters of Dolichos bean differed significantly due to foliar spray treatments of different organic nutrients. Among the different foliar treatments, seaweed extract spray @ 5% recorded significant highest plant height (96.13 cm), number of leaves plant<sup>-1</sup> (72.14), total chlorophyll in fresh leaf tissue (2.57 mg g<sup>-1</sup>), Leaf area (20.00 cm<sup>2</sup>), leaf area index (8.02), Days to first flowering (27.01 DAS), followed by T<sub>7</sub>. INM + Vermiwash @ 1:7.5 dilution and T<sub>10</sub>. - INM + Seaweed extract @ 2.5% foliar spray.

**Keywords:** Plant growth regulators, Dolichos bean.

## INTRODUCTION

Dolichos bean (*Lablab purpureus* (L) Sweet). belongs to the family leguminosae, is one of the important vegetable valued for its proteins, minerals, and energy. The crop is gaining popularity among vegetable growers due to higher remuneration and steady market demand. However, poor productivity level with traditional practices impairs wider acceptability of the crop. There is a tremendous scope to increase the current productivity level by adopting innovative practices. Different treatments of plant growth regulators were found effective in increasing the growth parameters through enhanced biomass production and translocation of assimilates toward developing sink. In addition to organic regulators like Seaweed (Sivasankari *et al.*, 2006), Vermiwash (Kaur *et al.*, 2015), Panchagavya (Swaminathan *et al.*, 2007), Effective micro organism (Chaudhary *et al.*, 2006) and Diammonium Phosphate (Kuldeep Singh *et al.*, 2015) are also emerging as plant booster for improving the physiological efficiency of the crop. In light of the fact, the present experiment was

undertaken to study the influences of certain plant growth regulators on growth attributes of Dolichos bean and to identify the most suitable combination of organic nutrients is along with INM.

## MATERIALS AND METHODS

The study was conducted during 2017 at farmer field in Varagoorpetai village, Cuddalore, Tamilnadu (  $11^{\circ} 24'$  N latitude,  $70^{\circ}41'$  E longitude and  $\pm 5.79$  m MSL above mean sea level). The soil was sandy loamy in texture, alkaline in reaction (pH 5.74) low in available nitrogen ( $53.0 \text{ kg ha}^{-1}$ ), higher in available phosphorus ( $31.0 \text{ kg P}_2\text{O}_5/\text{ha}$ ) and higher in potassium ( $145.0 \text{ kg K}_2\text{O}/\text{ha}$ ) respectively. The dolichos bean (cv. Co (Gb)-14 ) seeds were sown during mid - December in  $30 \times 60$  cm plant spacing. Five different growth regulators namely Diammonium Phosphate (2%), Panchagavya (3 and 4%), Vermiwash (1:5 and 1:7.5 dilution), Effective micro organism (1:1000 and 2.5: 1000 dilution). Seaweed extract (2.5 and 5%) along with control (distilled water) thus eleven treatments were laid out in Randomized Block Design with three replications. The experiment field was supplied with well rotten farmyard manure ( $20 \text{ t ha}^{-1}$ ) along with recommended dose of fertilizers NP ( $30: 50 \text{ kg ha}^{-1}$ ) +  $5 \text{ t VC ha}^{-1}$  + BF each  $2.5 \text{ kg ha}^{-1}$  to the plots and the growth regulators were sprayed at 30, 60 and 75 days after planting. The observations were recorded on five randomly selected plants from each plot on different growth characters (Table 1). The data was analyzed by adopting the standard procedure of Panse and Sukhatme (1985) and using AGRISTAT software. Wherever, the results were found significant, critical differences (CD) were computed at 5 percent level of probability to draw statistical conclusions.

### Growth parameters

The result revealed that the growth characters were significantly influence (Table I) as a result of foliar application of growth substances. Significantly higher plant height (96.13cm), number of leaves  $\text{plant}^{-1}$ (72.14) and total chlorophyll in fresh leaf tissue ( $2.75 \text{ mg g}^{-1}$ ) were observed treatment  $T_{11}$  (INM + Seaweed extract @ 5%) was applied to crop. Seaweed is involved in both cell division and cell elongation and can stimulate plant tissue resulting in enhanced vegetative growth (Rathore *et al.* 2009). The maximum leaf area ( $20.00 \text{ cm}^2$ ) and LAI (8.02) were registered by the same treatment ( $T_{11}$ ). However, both were closely followed by the treatments  $T_7$  (INM + Vermiwash @ 1:7.5 dilution) and  $T_{10}$  (INM + Seaweed extract @ 2.5%). Higher chlorophyll in leaves might have increased the photosynthesis rate and activated the leaf growth and subsequently the leaf area and LAI of the plant. The early flowering (27.01 days) in the treatment  $T_{11}$  (INM + Vermiwash @ 1:7.5 dilution) treated plants might be due to early completion of vegetative growth and better nourishment of plants. Kocira *et al.*, (2013) also reported increased vegetative growth, higher biomass production and early flowering of *Phaseolus vulgaris L.* with seaweed treatment.

Foliar spray of in EM, Panchagavya, and DAP emerged as second and third best option for most of the growth characters. Considering the growth characters, it may be concluded that for high growth characters plant height, number of leaves, chlorophyll content, leaf area and early flowering of coastal regain Dolichos bean application of INM + Seaweed extract @ 5% foliar spray at 30, 60 and 75 days of planting may be practiced for coastal region of Tamilnadu.

**Table 1: Effect of different organic growth regulators on growth parameters of dolichous bean**

Treatments	Plant height (cm)	No leaves /pant	Chlorophyll content fresh leaf tissue (mg g <sup>-1</sup> )	Leaf area (cm <sup>2</sup> )	LAI	Days to flowering (DAS)
T <sub>1</sub> - RDF (control)	54.51	53.86	2.01	17.78	5.32	36.67
T <sub>2</sub> - INM (Without foliar application)	54.06	56.73	2.11	18.06	5.69	34.74
T <sub>3</sub> - INM + DAP @ 2%	63.06	58.89	2.12	18.34	6.00	33.76
T <sub>4</sub> - INM + Panchagavya @ 3%	69.96	61.07	2.16	18.62	6.32	32.78
T <sub>5</sub> - INM + Panchagavya @ 4%	75.92	63.25	2.14	18.89	6.64	31.84
T <sub>6</sub> - INM + Vermiwash @ 1:5 dilution	83.57	67.62	2.19	19.44	7.30	29.73
T <sub>7</sub> - INM + Vermiwash @ 1:7.5 dilution	90.23	69.9	2.62	19.72	7.66	28.81
T <sub>8</sub> - INM + Effective Micro organism @ 1:1000 dilution	79.74	65.44	2.16	19.16	6.97	29.79
T <sub>9</sub> - INM + Effective Micro organism @ 2.5:1000 dilution	86.43	67.71	2.2	19.45	7.32	28.87
T <sub>10</sub> - INM + Seaweed extract @ 2.5% foliar spray	92.35	69.93	2.63	19.73	7.67	27.93
T <sub>11</sub> - INM + Seaweed extract @ 5% foliar spray	96.13	72.14	2.75	20.00	8.02	27.01
<b>SED</b>	<b>1.49</b>	<b>0.87</b>	<b>0.05</b>	<b>0.11</b>	<b>0.14</b>	<b>0.36</b>
<b>CD</b>	<b>3.10</b>	<b>1.81</b>	<b>0.10</b>	<b>0.22</b>	<b>0.29</b>	<b>0.75</b>

INM: NP (30: 50 kg ha<sup>-1</sup>) + VC @ 5 t ha<sup>-1</sup> + BF 2.5 kg ha<sup>-1</sup>

## REFERENCE

- Chaudhary MS, Iqupa M (2006) Soil fertility improvements with EM for vegetables crops. EM database, EM technology network, Inc.
- Kaur P., Bhardwaj M. and Babbar I. (2015) Effect of Vermicompost and Vermiwash on Growth of Vegetables. Res. J. Animal, Veterinary and Fishery Sci., Vol. 3 (4), 9-12,
- Kocira A, Kornas R, Kocira S. 2013. Effect assessment of Kelpak sl on the Bean yield. Journal of Central European Agriculture, 14 (2), p.67- 76.
- Kuldeep Singh, R.S. Manohar, Rakesh Choudhary, A.K. Yadav and A. Sangwan (2015). Response of different sources and levels of phosphorus on yield, nutrient uptake and net returns on mungbean under rainfed condition. Agric. Sci. Digest., 35 (4): 263-268
- Panse, V.G. and Sukhatme, P.V. (1985) Statistical Methods for Agricultural Workers. Indian Council of Agricultural Research Publication 87-89.
- Rathore, S.S. Chaudhary, D.R. Boricha, G.N. Ghosh A. Bhatt, B.P. Zodape, S.T. Patolia J.S. (2009) Effect of seaweed extract on the growth, yield and nutrient uptake of soybean (Glycine max) under rainfed conditions. South African Journal of Botany 75 :351–355.
- Sivasankari, S., Venkatesalu, V., Anantharaj, M., Chandrasekaran, M., 2006. Effect of seaweed extracts on the growth and biochemical constituents of Vigna sinensis. Bioresource Technology 97, 1745–1751.
- Swaminathan, C., Swaminathan, V., and Vijayalakshmi, V. (2007). Panchagavya Boon to Organic Farming. International Book Distributing Co., India.